arXiv:1406.6991v2 [math.NT] 29 Jul 2014

Asymptotic behavior of a series of Euler's totient function $\varphi(k)$ times the index of $1/k$ in a Farey sequence

R. Tomás *
CERN, CH 1211 Geneva 23, Switzerland

July 30, 2014

Abstract

Motivated by studies in accelerator physics this paper computes the asymptotic behavior of the series $\sum_{k=1}^{N} \varphi(k) I_N\left(\frac{1}{k}\right)$, where $\varphi(k)$ is Euler's totient function and $I_N\left(\frac{1}{k}\right)$ is the position that $1/k$ occupies in the Farey sequence of order N. To this end an exact formula for $I_N\left(\frac{1}{k}\right)$ is derived when all integers in $\left[2, \left\lceil \frac{N}{k} \right\rceil\right]$ are divisors of N.

1 Results

Let $I_N\left(\frac{1}{k}\right)$, with $k \in \mathbb{N}$ and $k \leq N$, be the position that $1/k$ occupies in the Farey sequence of order N. Some useful facts follow:

$$I_N\left(\frac{0}{1}\right) = 1 \,, \tag{1}$$

$$I_N\left(\frac{1}{1}\right) = |F_N| \,, \tag{2}$$

$$I_N\left(\frac{1}{k}\right) = 1 + \sum_{j=k}^{N} \phi\left(j; \left[1, \left\lfloor \frac{j}{k} \right\rfloor\right]\right) \,, \tag{3}$$

$$I_N\left(\frac{1}{k}\right) = 1 + |F_N| - I_N\left(\frac{k-1}{k}\right) \,, \tag{4}$$

*rogelio.tomas@cern.ch

where $|F_N|$ stands for the cardinality of the Farey sequence of order N and Eq. (3) is found at [1] Remark 7.10 (there fractions are indexed starting with 0); $\phi(n;[\cdot,\cdot])$ is defined as the number of elements from an interval of integers that are relatively prime to n. Consequently $\phi(n;[1,n]) \equiv \varphi(n)$ and $|F_N| = 1 + \sum_{j=1}^{N} \phi(j;[1,j])$. Efficient algorithms for the computation of $I_N(x)$ were recently developed in [2].

Theorem 1.1. *Given $k \in \mathbb{N}$ such that $k \leq N$, then*

$$I_N\left(\frac{1}{k}\right) \leq \frac{N^2+N}{2k} \ . \tag{5}$$

Proof. In Eq. (3) the set $\left[1, \left\lfloor \frac{j}{k} \right\rfloor\right]$ has a maximum of j/k elements with j running between k and N. Therefore ϕ tests a maximum of

$$\sum_{j=k}^{N} \frac{j}{k} \quad = \quad \frac{(N-k+1)(N+k)}{2k} = \frac{N^2-k^2+N+k}{2k} \tag{6}$$

elements. □

Let the subsequence of F_N, $F_N^{1/a,\,1/b}$ be defined as all the fractions of F_N in $[1/a,\ 1/b]$ with $1 \leq b \leq a \leq N$.

Theorem 1.2. *If N is a multiple of i and $i+1$ there is a bijective and order-preserving map between F_i and $F_N^{1/q,\,1/(q-1)}$, with q being an integer fulfilling $N/(i+1) < q \leq N/i$, given by*

$$F_i \to F_N^{1/q,\,1/(q-1)} \ , \qquad \frac{h}{k} \mapsto \frac{k}{kq-h} \ . \tag{7}$$

Proof. The demonstration is given in two steps. The first step is shown in Fig. 1, where by construction it is clear that all Farey fractions in $[1/q,\, 1/(q-1)]$ at any order are connected to the Farey fractions in $[0/1,\, 1/1]$ by the application in Eq. (7).

A second step is needed to show that the application is bijective between the sets F_i and $F_N^{1/q,\,1/(q-1)}$. The fraction h/k belongs to F_i if $k \leq i$, and similarly $k/(kq-h)$ belongs to $F_N^{1/q,\,1/(q-1)}$ if $(kq-h) \leq N$. Since N is a multiple of i and $i+1$ and that $N/(i+1) < q \leq N/i$, the largest value $(kq-h)$ takes is $\left(k\frac{N}{i} - h\right)$. Therefore $k/(kq-h)$ belongs to $F_N^{1/q,\,1/(q-1)}$ if $k \leq i$, i.e., if h/k belongs to F_i.

In the opposite direction, if $(kq-h) \leq N$ the largest possible k is obtained by inserting the smallest q, which is $N/(i+1)+1$, yielding

$$k\left(\frac{N}{i+1}+1\right) - h \leq N \ . \tag{8}$$



Figure 1: Application between the Farey fractions in $[1/q, 1/(q-1)]$ and $[0,1]$ demonstrated by using that the next Farey fraction appearing between h/k and h'/k' is $(h+h')/(k+k')$. By applying this rule independently to $1/q$ and $1/(q-1)$ (top) and $0/1$ and $1/1$ (bottom) the map of Eq. (7) is apparent. The second and the last to the last terms of F_i (bottom) and $F_N^{1/q,\,1/(q-1)}$ (top) are also shown. They can be computed using Corollary 3.2 of [3].

If $k = h$ this corresponds to the trivial case $h/k = 1/1$, which clearly satisfies Eq. (1). Else, $k > h$ and Eq. (8) becomes

$$k\frac{N}{i+1} < N \ , \tag{9}$$

$$k \leq i \ , \tag{10}$$

concluding that if $k/(kq-h)$ belongs to $F_N^{1/q,\,1/(q-1)}$, h/k belongs to F_i. □

Note that in the case $q = 2$ the map in Eq. (7) can be viewed as a map from [3] Lemma 1.1 that reflects a Farey sequence to the right half sequence of a Farey subsequence arising in the combinatorics of finite sets.

Theorem 1.3. *Let $N/(i+1) \leq k \leq N/i$ and N be a multiple of all integers in $[2, i]$, then*

$$I_N\left(\frac{1}{k}\right) = 2 + N\sum_{j=1}^{i}\frac{\varphi(j)}{j} - k\Phi(i) \ , \tag{11}$$

where $\Phi(i)$ is the totient summatory function, $\Phi(i) \equiv \sum_{j=1}^{i}\varphi(j) \equiv |F_i| - 1$.

Proof. Express $I_N(1/(N/i))$ as the sum of the cardinalities of all subsequences of the form $F_N^{1/q,1/(q-1)}$ such that $q > N/i$,

$$I_N\left(\frac{1}{N/i}\right) = 2 + \sum_{q=N/i+1}^{N}\left(\left|F_N^{1/q,1/(q-1)}\right| - 1\right) \ . \tag{12}$$

By virtue of Theorem 1.2, $\left|F_N^{1/q,1/(q-1)}\right| = |F_i|$ when $N/(i+1) < q \leq N/i$ and Eq. (12) is directly re-written as

$$I_N\left(\frac{1}{N/i}\right) = 2 + N\sum_{j=1}^{i-1}\frac{\Phi(j)}{j(j+1)} \ . \tag{13}$$

$I_N(1/k)$ is computed by adding to the expression above the cardinality of the remaining subsequences between N/i and k, yielding

$$I_N\left(\frac{1}{k}\right) = \Phi(i)\left(\frac{N}{i} - k\right) + I_N\left(\frac{1}{N/i}\right) \ . \tag{14}$$

After some algebra Eq. (11) is obtained. □

Theorem 1.4.

$$\sum_{k=1}^{N}\varphi(k)I_N(1/k) = \frac{N^3}{6\zeta(3)} + O\left(\frac{N^3}{\log N}\right) \tag{15}$$

Proof. Let $N = \mathrm{lcm}(2,3,4,...,i_{max})$ be the least common multiple of the first i_{max} numbers. The summation in the left hand side of Eq. (15) is split into two parts, the first part being for $k \leq N/i_{max}$. Equation (5) is used to give an upper bound to the summation corresponding to this first part. In the second part, $k > N/i_{max}$, Eq. (11) is used. The two contributions are given by

$$\begin{aligned}
\sum_{k=1}^{N/i_{max}} \varphi(k) I_N(1/k) &\leq (N^2+N) \sum_{k=1}^{N/i_{max}} \frac{\varphi(k)}{2k} , \qquad (16)\\
\sum_{k>N/i_{max}}^{N} \varphi(k) I_N(1/k) &= \sum_{i=1}^{i_{max}-1} \sum_{k=\frac{N}{i+1}+1}^{N/i} \varphi(k) \left(2 + N \sum_{j=1}^{i} \frac{\varphi(j)}{j} - k\Phi(i) \right)
\end{aligned}$$

After some algebra using the following relations,

$$\begin{aligned}
\sum_{k=1}^{N} \varphi(k) &= \frac{3}{\pi^2} N^2 + O(N \log N) , \qquad (17)\\
\sum_{k=1}^{N} \frac{\varphi(k)}{k} &= \frac{6}{\pi^2} N + O((\log N)^{2/3} (\log\log N)^{4/3}) , \qquad (18)\\
\sum_{k=1}^{N} \varphi(k) k &= \frac{2}{\pi^2} N^3 + O(N^2) , \qquad (19)\\
\sum_{k=1}^{\infty} \frac{\varphi(k)}{k^3} &= \frac{\zeta(2)}{\zeta(3)} = \frac{\pi^2}{6\zeta(3)} , \qquad (20)
\end{aligned}$$

one obtains

$$\begin{aligned}
\sum_{k=1}^{N/i_{max}} \varphi(k) I_N(1/k) &\leq \frac{3}{\pi^2 i_{max}} (N^3 + N^2) , \qquad (21)\\
\sum_{k>N/i_{max}}^{N} \varphi(k) I_N(1/k) &= \frac{N^3}{\pi^2} \sum_{i=1}^{i_{max}} \frac{\varphi(i)}{i^3} + O(N^2 \log N) . \qquad (22)
\end{aligned}$$

When taking the limit $i_{max} \to \infty$, N tends to $e^{i_{max}}$ and, inversely, i_{max} tends to $\ln(N)$. Equations (21) and (22) become

$$\begin{aligned}
\sum_{k=1}^{N/i_{max}} \varphi(k) I_N(1/k) &\leq \frac{3}{\pi^2} \frac{N^3 + N^2}{\ln(N)} , \qquad (23)\\
\sum_{k>N/i_{max}}^{N} \varphi(k) I_N(1/k) &= \frac{N^3}{6\zeta(3)} + O(N^2 \log N) . \qquad (24)
\end{aligned}$$

Combining these two last equations Eq. (15) is obtained. □

2 Discussion

Let T_N^D be the number of linear, integral and irreducible polynomials of dimension D of the form

$$\sum_{i=1}^{D} a_i x_i - a_{D+1} \ , \tag{25}$$

such that $\sum_{i=1}^{D} |a_i| \leq N$ and having at least one root in the unitary $D-$cube. $D = 0$ corresponds to the null polynomial 0. $D = 1$ corresponds to Farey sequences. The result of the previous section serves to compute T_N^2, see [4, 5]. The asymptotic behavior in N of T_N^D, for $D = 0, 1, 2$ follow,

$$T_N^0 = 1 \ , \tag{26}$$

$$T_N^1 = \frac{3}{\pi^2} N^2 + O(N \log N) \ , \tag{27}$$

$$T_N^2 = \frac{2N^3}{3\zeta(3)} + O\left(\frac{N^3}{\log N}\right) \ . \tag{28}$$

3 Acknowledgments

Andrey Matveev is warmly thanked for making very useful suggestions on the manuscript.

References

[1] Andrey O. Matveev, *Relative blocking in posets*, Journal of Combinatorial Optimization, **13** (2007) 379–403. *Corrigendum*: `arXiv:math/0411026`

[2] J. Pawlewicz and M. Pătraşcu, *Order Statistics in the Farey Sequences in Sublinear Time and Counting Primitive Lattice Points in Polygons*, Algorithmica, Volume **55**, Issue 2, pp 271-282, 2009.

[3] Andrey O. Matveev, *Neighboring fractions in Farey subsequences*, `arXiv:math/0801.1981`

[4] R. Tomás, *From Farey sequences to resonance diagrams* Phys. Rev. ST Accel. Beams **17**, 014001 (2014).

[5] H. Brown and K. Mahler, *A Generalization of Farey Sequences: Some Exploration Via the Computer*, Journal of Number Theory, Vol. **3**, 364, August 1971.